

附录 1 数据来源

Appendix 1 Data sources of 70 study sites

| | | | |
|----|--------------------------------|----|-------------------------------|
| 1 | Sundarapandian and Swamy, 1998 | 36 | Zhou et al., 2014 |
| 2 | Jessy et al., 2013 | 37 | Wang et al., 2012 |
| 3 | Jessy et al., 2013 | 38 | Zhou et al., 2014 |
| 4 | Visalakshi, 1994 | 39 | Li et al., 2013 |
| 5 | 伍恩华和刘强, 2013 | 40 | 梅莉等, 2006 |
| 6 | Xu et al., 2013 | 41 | 王永吉等, 2014 |
| 7 | 郝瑞, 2007 | 42 | Fukuzawa et al., 2013 |
| 8 | Arunachalam et al., 1996 | 43 | 耿鹏飞, 2015 |
| 9 | 郑金兴等, 2012 | 44 | Yang et al., 2010 |
| 10 | 廖利平等, 1995 | 45 | Adamek et al., 2011 |
| 11 | 曹乐, 2014 | 46 | Parrotta and Lodge, 1991 |
| 12 | 李树战等, 2008 | 47 | Carter et al., 2004 |
| 13 | 何永涛等, 2009 | 48 | Farrish, 1991 |
| 14 | 曾凡荣等, 2008 | 49 | Hart et al., 2005 |
| 15 | 吴勇, 2011 | 50 | Powell and Day, 1991 |
| 16 | 董慧霞和李贤伟, 2016 | 51 | Dress and Boerner, 2001 |
| 17 | 王力等, 2013 | 52 | Wilczynski and Pichett, 1993 |
| 18 | 刘波, 2008 | 53 | Zhou et al., 2011 |
| 19 | Garkoti, 2011 | 54 | Son and Hwang, 2003 |
| 20 | 王哲等, 2014 | 55 | Burke and Raynal, 1994 |
| 21 | 王哲等, 2014 | 56 | Santantonio and Hermann, 1985 |
| 22 | Jiang et al., 2016 | 57 | Andersen et al., 2008 |
| 23 | Singh and Singh, 2016 | 58 | Cronan, 2003 |
| 24 | 陈金林等, 1999 | 59 | Keyes and Grier, 1981 |
| 25 | Van Do et al., 2015 | 60 | Brassard et al., 2013 |
| 26 | Ma et al., 2013 | 61 | Montagnoli et al., 2012 |
| 27 | Tawa and Takeda, 2016 | 62 | Kummerow et al., 1990 |
| 28 | 李盼盼, 2012 | 63 | Bakker, 1999 |
| 29 | Van Do et al., 2016 | 64 | Meinen et al., 2009 |
| 30 | Konôpka et al., 2006, 2007 | 65 | Berhongaray et al., 2015 |
| 31 | Son and Hwang, 2003 | 66 | Ostonen et al., 2005 |
| 32 | 闫美芳等, 2015 | 67 | Makkonen and Helmisaari, 1999 |
| 33 | Xiao et al., 2008 | 68 | Silver et al., 2005 |
| 34 | Pei et al., 2012 | 69 | Lima et al., 2010 |
| 35 | Wang et al., 2014 | 70 | Schroth and Zech, 1995 |

曹乐 (2014). 凤阳山不同林分类型细根生物量及周转率研究. 硕士学位论文. 南京: 南京林业大学.

陈金林, 许新健, 姜志林, 张武兆, 张井义, 贾永正 (1999). 空青山次生栎林细根周转. 南京林业大学学报 **23**, 6–10.

- 董慧霞, 李贤伟 (2016). 不同草本层三倍体毛白杨细根及草根的空间分布及季节变化. 河南农业科学 **45**, 108–113.
- 耿鹏飞 (2015). 小兴安岭 4 种林型细根生物量时空格局及其养分分布. 硕士学位论文. 哈尔滨: 东北林业大学.
- 郝瑞 (2007). 沿海沙地竹林细根生态系统特性研究. 硕士学位论文. 福州: 福建农林大学.
- 何永涛, 石培礼, 张宪洲, 钟志明, 徐玲玲, 张东秋 (2009). 拉萨河谷杨树人工林细根的生产力及其周转. 生态学报 **29**, 2877–2883.
- 李盼盼 (2012). 杨树人工林细根的空间分布特征及其季节动态. 硕士学位论文. 泰安: 山东农业大学.
- 李树战, 田大伦, 王光军, 闫文德, 朱凡 (2008). 樟树人工林细根生物量的时空动态. 中南林业科技大学学报 **28**(6), 51–55.
- 廖利平, 陈楚莹, 张家武, 高洪 (1995). 杉木、火力楠纯林及混交林细根周转的研究. 应用生态学报 **6**, 7–10.
- 刘波 (2008). 亚热带常绿阔叶林细根生物量及根系呼吸的研究. 硕士学位论文. 合肥: 安徽农业大学.
- 梅莉, 韩有志, 于水强, 史建伟, 王政权 (2006). 水曲柳人工林细根季节动态及其影响因素. 林业科学 **42**(9), 7–12.
- 王力, 吴福金, 邓文鑫, 余学明, 吴磊 (2013). 马尾松混交林细根生物量研究. 安徽农业大学学报 **40**, 225–232.
- 王永吉, 赵学春, 来利明, 朱林海, 王健健, 周继华, 姜联合, 马远见, 赵春强, 郑元润 (2014). 沙枣人工群落细根生物量和周转过程. 干旱区地理 **37**, 548–554.
- 王哲, 冯春, 晋曦, 马严, 傅松玲, 陈汉, 杜勤智 (2014). 不同封育年限下石灰岩山地植被细根生物量及其动态. 广东农业科学 **41**(3), 170–174.
- 伍恩华, 刘强 (2013). 海南岛北部木麻黄林凋落物与土壤养分、细根生物量的关系. 广西植物 **33**, 488–495.
- 吴勇 (2011). 台湾栲木林草复合细根特性研究. 博士论文. 雅安: 四川农业大学.
- 闫美芳, 王璐, 郭楠, 时红 (2015). 黄土高原杨树人工林的细根生物量与碳储量研究. 中国农学通报 **31**(35), 146–151.
- 曾凡荣, 施家月, 阎恩荣, 张仁良, 王希华 (2008). 天童常绿阔叶林次生演替过程中细根的生物量动态. 华东师范大学学报(自然科学版) **(6)**, 56–62.
- 郑金兴, 陈光水, 高人, 杨智杰 (2012). 楠木人工林细根生物量季节动态及周转. 三明学院学报 **29**(6), 89–94.
- Adamek M, Corre MD, Hölscher D** (2011). Responses of fine roots to experimental nitrogen addition in a tropical lower montane rain forest, Panama. *J Trop Ecol* **27**, 73–81.
- Andersen CP, Phillips DL, Rygiewicz PT, Storm MJ** (2008). Fine root growth and mortality in different-aged ponderosa pine stands. *Can J For Res* **38**, 1797–1806.

- Arunachalam A, Pandey HN, Tripathi RS, Maithani K** (1996). Biomass and production of fine and coarse roots during regrowth of a disturbed subtropical humid forest in north-east India. *Vegetatio* **123**, 73–80.
- Bakker MR** (1999). The effect of lime and gypsum applications on a sessile oak (*Quercus petraea* (M.) Liebl.) stand at La Croix-Scaille (French Ardennes) II. Fine root dynamics. *Plant Soil* **206**, 109–121.
- Berhongaray G, Verlinden MS, Broeckx LS, Ceulemans R** (2015). Changes in belowground biomass after coppice in two *Populus* genotypes. *For Ecol Manage* **337**, 1–10.
- Brassard BW, Chen HYH, Cavard X, Laganière J, Reich PB, Bergeron Y, Parè D, Yuan ZY** (2013). Tree species diversity increases fine root productivity through increased soil volume filling. *J Ecol* **101**, 210–219.
- Burke MK, Raynal DJ** (1994). Fine root growth phenology, production, and turnover in a northern hardwood forest ecosystem. *Plant Soil* **162**, 135–146.
- Carter DC, Hendricks JJ, Mitchell RJ, Pecot SD** (2004). Fine root carbon allocation and fates in longleaf pine forests. *Forest Science* **50**, 177–187.
- Cronan CS** (2003). Belowground biomass, production, and carbon cycling in mature Norway spruce, Maine, USA. *Can J For Res* **33**, 339–350.
- Dress WJ, Boerner REJ** (2001). Root dynamics of southern Ohio oak-hickory forests: influences of prescribed fire and landscape position. *Can J For Res* **31**, 644–653.
- Farrish KW** (1991). Spatial and temporal fine-root distribution in Three Louisiana forest soils. *Soil Sci Soc Am J* **55**, 1752–1757.
- Fukuzawa K, Shibata H, Takagi K, Satoh F, Koike T, Sasa K** (2013). Temporal variation in fine-root biomass, production and mortality in a cool temperate forest covered with dense understory vegetation in northern Japan. *For Ecol Manage* **310**, 700–710.
- Garkoti SC** (2011). Fine root dynamics in three central Himalayan high elevation forests ranging from closed canopied to open-canopied treeline vegetation. *J For Res* **16**, 136–143.
- Hart SC, Classen AT, Wright RJ** (2005). Long-term interval burning alters fine root and mycorrhizal dynamics in a ponderosa pine forest. *J Appl Ecol* **42**, 752–761.
- Jessy MD, Prasannakumari P, Abraham J** (2013). Carbon and nutrient cycling through fine roots in rubber (*Hevea brasiliensis*) plantations in India. *Exp Agric* **49**, 556–573.
- Jiang H, Bai YY, Du HY, Hu Y, Rao YF, Chen C, Cai YL** (2016). The spatial and seasonal variation characteristics of fine roots in different plant configuration modes in new reclamation saline soil of humid climate in China. *Ecol Eng* **86**, 231–238.
- Keyes MR, Grier CC** (1981). Above- and below-ground net production in 40-year-old Douglas-fir stands on low and high productivity sites. *Can J For Res* **11**, 599–605.

- Konôpka B, Noguchi K, Sakata T, Takahashi M, Konôpková Z** (2006). Fine root dynamics in a Japanese cedar (*Cryptomeria japonica*) plantation throughout the growing season. *For Ecol Manage* **225**, 278–286.
- Konôpka B, Noguchi K, Sakata T, Takahashi M, Konôpková Z** (2007). Effects of simulated drought stress on the fine roots of Japanese cedar (*Cryptomeria japonica*) in a plantation forest on the Kanto Plain, eastern Japan. *J For Res* **12**, 143–151.
- Kummerow J, Kummerow M, Trabaud L** (1990). Root biomass, root distribution and the fine-root growth dynamics of *Quercus coccifera* L. in the garrigue of southern France. *Vegetatio* **87**, 37–44.
- Li XF, Zhu J, Lange H, Han SJ** (2013). A modified ingrowth core method for measuring fine root production, mortality and decomposition in forests. *Tree Physiol* **33**, 18–25.
- Lima TTS, Miranda ISM, Vasconcelos SS** (2010). Effects of water and nutrient availability on fine root growth in eastern Amazonian forest regrowth, Brazil. *New Phytol* **187**, 622–630.
- Ma C, Zhang WH, Wu M, Xue YQ, Ma LW, Zhou JY** (2013). Effect of aboveground intervention on fine root mass, production, and turnover rate in a Chinese cork oak (*Quercus variabilis* Blume) forest. *Plant Soil* **368**, 201–214.
- Makkonen K, Helmisaari HS** (1999). Assessing fine-root biomass and production in a Scots pine stand—comparison of soil core and root ingrowth core methods. *Plant Soil* **210**, 43–50.
- Meinen C, Hertel D, Leuschner C** (2009). Root growth and recovery in Temperate Broad-Leaved Forest stands differing in tree species diversity. *Ecosystems* **12**, 1103–1116.
- Montagnoli A, Terzaghi M, Di Iorio A, Scippa GS, Chiatante D** (2012). Fine-root morphological and growth traits in a Turkey-oak stand in relation to seasonal changes in soil moisture in the Southern Apennines, Italy. *Ecol Res* **27**, 1015–1025.
- Ostonen I, Löhmus K, Pajuste K** (2005). Fine root biomass, production and its proportion of NPP in a fertile middle-aged Norway spruce forest: comparison of soil core and ingrowth core methods. *For Ecol Manage* **212**, 264–277.
- Parrotta JA, Lodge DJ** (1991). Fine root dynamics in a subtropical wet forest following hurricane disturbance in Puerto Rico. *Biotropica* **23**, 343–347.
- Pei ZQ, Xiao CW, Dong D, Zhang SR** (2012). Comparison of the fine root dynamics of *Populus euphratica* forests in different habitats in the lower reaches of the Tarim River in Xinjiang, China, during the growing season. *J For Res* **17**, 343–351.
- Powell SW, Day Jr FP** (1991). Root production in four communities in the great dismal swamp. *Am J Bot* **78**, 288–297.

- Santantonio D, Hermann RK** (1985). Standing crop, production, and turnover of fine roots on dry, moderate, and wet sites of mature Douglas-fir in western Oregon. *Ann For Sci* **42**, 113–142.
- Schroth G, Zech W** (1995). Above- and below-ground biomass dynamics in a sole cropping and an alley cropping system with *Gliricidia sepium* in the semi-deciduous rainforest zone of west Africa. *Agroforestry Systems* **31**, 181–198.
- Silver WL, Thompson AW, McGroddy ME, Varner RK, Dias JD, Silva H, Crill PM, Keller M** (2005). Fine root dynamics and trace gas fluxes in two lowland tropical forest soils. *Glob Chang Biol* **11**, 292–306.
- Singh P, Singh B** (2016). Biomass and nitrogen dynamics of fine roots of poplar under differential N and P levels in an agroforestry system in Punjab. *Trop Ecol* **57**, 143–152.
- Son Y, Hwang JH** (2003). Fine root biomass, production and turnover in a fertilized *Larix leptolepis* plantation in central Korea. *Ecol Res* **18**, 339–346.
- Sundarapandian S, Swamy P** (1998). Variation in fine-root biomass and net primary productivity due to conversion of tropical forests into plantation crops and agroecosystems. In: Box Jr JE, ed. *Root Demographics and their Efficiencies in Sustainable Agriculture, Grasslands and Forest Ecosystems*. Dordrecht: Springer. pp. 369–382.
- Tawa Y, Takeda H** (2016). Fine root dynamics in organic and mineral soil layers of *Cryptomeria japonica* D. Don plantation. *J For Res* **21**, 67–72.
- Van Do T, Sato T, Koza O** (2016). A new approach for estimating fine root production in forests: a combination of ingrowth core and scanner. *Trees* **30**, 545–554.
- Van Do T, Sato T, Saito S, Koza O** (2015). Fine-root production and litterfall: main contributions to net primary production in an old-growth evergreen broad-leaved forest in southwestern Japan. *Ecol Res* **30**, 921–930.
- Visalakshi N** (1994). Fine root dynamics in two tropical dry evergreen forests in southern India. *J Biosci* **19**, 103–116.
- Wang CG, Han SJ, Zhou YM, Yan CF, Cheng XB, Zheng XB, Li MH** (2012). Responses of fine roots and soil N availability to short-term nitrogen fertilization in a broad-leaved Korean pine mixed forest in Northeastern China. *PLoS One* **7**, e31042.
- Wang XY, Ma LY, Jia ZK, Jia LM** (2014). Root inclusion net method: novel approach to determine fine root production and turnover in *Larix principis-rupprechtii* Mayr plantation in North China. *Turk J Agric For* **38**, 388–398.
- Wilczynski CJ, Pichett STA** (1993). Fine root biomass within experimental canopy gaps: evidence for a below-ground gap. *J Veg Sci* **4**, 571–574.
- Xiao CW, Sang WG, Wang RZ** (2008). Fine root dynamics and turnover rate in an Asia white birch forest of Donglingshan Mountain, China. *For Ecol Manage* **255**, 765–773.

- Xu WQ, Liu JX, Liu XZ, Li K, Zhang DQ, Yan JH** (2013). Fine root production, turnover, and decomposition in a fast-growth *Eucalyptus urophylla* plantation in southern China. *J Soils Sediments* **13**, 1150–1160.
- Yang LY, Wu ST, Zhang LB** (2010). Fine root biomass dynamics and carbon storage along a successional gradient in Changbai Mountains, China. *Forestry* **83**, 379–387.
- Zhou Y, Su JQ, Janssens IA, Zhou GS, Xiao CW** (2014). Fine root and litterfall dynamics of three Korean pine (*Pinus koraiensis*) forests along an altitudinal gradient. *Plant Soil* **374**, 19–32.
- Zhou YM, Tang J, Melillo JM, Butler S, Mohan JE** (2011). Root standing crop and chemistry after six years of soil warming in a temperate forest. *Tree Physiol* **31**, 707–717.